

Billing Codes, National Drug Codes, and Immunization Registry Codes for VAXELIS™ (Diphtheria and Tetanus Toxoids and Acellular Pertussis, Inactivated Poliovirus, Haemophilus b Conjugate and Hepatitis B Vaccine)

Included are lists of codes that may be relevant for VAXELIS and its administration.

This information is current as of November 2020. The information provided here is compiled from sources believed to be accurate, but Merck and Sanofi Pasteur make no representation that it is accurate. Consult the relevant manual and/or other guidelines for a description of each code to determine the appropriateness of a particular code and for information on additional codes. This information is subject to change. Merck and Sanofi Pasteur caution that payer coding requirements vary and can frequently change, so it is important to regularly check with each payer as to payer-specific requirements.

You are solely responsible for determining the appropriate codes and for any action you take in billing. The information provided here is not intended to be definitive or exhaustive, and is not intended to replace the guidance of a qualified professional advisor. Merck and Sanofi Pasteur and their agents make no warranties or guarantees, expressed or implied, concerning the accuracy or appropriateness of this information for your particular use given the frequent changes in public and private payer billing. The use of this information does not guarantee payment or that any payment received will cover your costs. Diagnosis codes should be selected only by a health care professional.

Indication

VAXELIS is a vaccine indicated for active immunization to prevent diphtheria, tetanus, pertussis, poliomyelitis, hepatitis B, and invasive disease due to *H. influenzae* type b. VAXELIS is approved for use as a 3-dose series in children 6 weeks through 4 years of age (prior to the 5th birthday).

Important Safety Information

- Do not administer VAXELIS to anyone with a history of severe allergic reaction to a previous dose of VAXELIS, any ingredient of VAXELIS, or any other diphtheria toxoid, tetanus toxoid, pertussis-containing vaccine, inactivated poliovirus vaccine, hepatitis B vaccine, or Hib vaccine.
- Do not administer VAXELIS to anyone with a history of encephalopathy within 7 days of a pertussis-containing vaccine with no other identifiable cause.
- Do not administer VAXELIS to anyone with a history of progressive neurologic disorder until a treatment regimen has been established and the condition has stabilized.
- Carefully consider benefits and risks before administering VAXELIS to persons with a history of: fever $\geq 40.5^{\circ}\text{C}$ ($\geq 105^{\circ}\text{F}$), hypotonic-hyporesponsive episode (HHE), or persistent, inconsolable crying lasting ≥ 3 hours within 48 hours after a previous pertussis-containing vaccine; and/or seizures within 3 days after a previous pertussis-containing vaccine.
- If Guillain-Barré syndrome occurred within 6 weeks of receipt of a prior vaccine containing tetanus toxoid, the risk for Guillain-Barré syndrome may be increased following VAXELIS.

Important Safety Information continued on pages 2 and 3.

VAXELIS™ (Diphtheria and Tetanus Toxoids and Acellular Pertussis, Inactivated Poliovirus, Haemophilus b Conjugate and Hepatitis B Vaccine)

Billing Codes

Common CPT® Codes ¹	Description
Vaccine product	
90697	Diphtheria, tetanus toxoids, acellular pertussis vaccine, inactivated poliovirus vaccine, <i>Haemophilus influenzae</i> type b PRP-OMP conjugate vaccine, and hepatitis B vaccine (DTaP-IPV-Hib-HepB), for intramuscular use
Administration of vaccine	
90460	Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified HCP; first or only component of each vaccine or toxoid administered (Bill 1 unit for the first component of VAXELIS)
and	
+90461	Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified HCP; each additional vaccine or toxoid component administered (list separately in addition to code for primary procedure) (Bill 5 units; 1 unit for each additional component within VAXELIS)
or	
90471	Immunization administration; 1 injected vaccine (single or combination vaccine/toxoid) (Assuming VAXELIS is administered first during child's visit, bill 1 unit)
+90472	Immunization administration; each additional injected vaccine (single or combination vaccine/toxoid) (Assuming VAXELIS is administered subsequent to other vaccine[s] during child's visit, bill 1 unit)
Common ICD-10-CM Code ²	Description
Z23 ^a	Encounter for immunization

^aZ23 is reported for all vaccine-related encounters for all vaccines given, and in addition to any health exam ICD-10-CM codes.¹

CPT, Current Procedural Terminology; HCP, health care provider; ICD-10-CM, International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Clinical Modification.

CPT is a registered trademark of the American Medical Association.

The use of this information does not guarantee payment or that payment received will cover your costs. Diagnosis codes should be selected only by a health care professional.

Important Safety Information (*continued*)

- Apnea following intramuscular vaccination has been observed in some infants born prematurely. Consider the individual infant's medical status and potential benefits and possible risks of intramuscular vaccination in deciding when to administer VAXELIS to an infant born prematurely.
- Vaccination with VAXELIS may not protect all individuals.

Important Safety Information continued on page 3.

National Drug Codes and Immunization Registry Codes

Supplied Packages	NDC Unit of Use	NDC Carton
10 single-dose vials	N463361024358 ML0.5	N463361024310 ML0.5
10 single-dose prefilled syringes	N463361024388 ML0.5	N463361024315 ML0.5
CVX Code ³	146	
MVX Code ³	PMC	

NDC, National Drug Code.

Please note: The NDCs above are the billable NDCs that appear on the carton and the unit of use, when NDC is required by the payer.

CVX Code, indicates the product used in a vaccination; MVX Code, indicates the manufacturer of a vaccine.

Codes for Administration⁴

If a vaccination is the only service provided, a visit should not be billed. If a vaccination is provided during a visit with other services, consult payer-specific requirements.

Important Safety Information (*continued*)

- The solicited adverse reactions 0-5 days following any dose were irritability ($\geq 55\%$), crying ($\geq 45\%$), injection site pain ($\geq 44\%$), somnolence ($\geq 40\%$), injection site erythema ($\geq 25\%$), decreased appetite ($\geq 23\%$), fever $\geq 38.0^{\circ}\text{C}$ ($\geq 19\%$), injection site swelling ($\geq 18\%$), and vomiting ($\geq 9\%$).
- The 3-dose immunization series consists of a 0.5 mL intramuscular injection, administered at 2, 4, and 6 months of age.
- A 3-dose series of VAXELIS does not constitute a primary immunization series against pertussis; an additional dose of pertussis-containing vaccine is needed to complete the primary series.

Before administering VAXELIS, please read the accompanying Prescribing Information. The Patient Information also is available.

References: **1.** American Academy of Pediatrics. Commonly administered pediatric vaccines. Page Updated August 1, 2019. Accessed October 15, 2020. https://www.aap.org/en-us/Documents/coding_vaccine_coding_table.pdf. **2.** Centers for Disease Control and Prevention. ICD-10-CM Tabular list of diseases and injuries. Page Last Modified March 31, 2020. Accessed October 15, 2020. <https://www.cms.gov/Medicare/Coding/ICD10/2020-ICD-10-CM>. **3.** Centers for Disease Control and Prevention. IIS: HL7 Standard code set: Mapping product names to CVX and MVX. Page Last Reviewed February 12, 2020. Accessed October 15, 2020. <https://www2a.cdc.gov/vaccines/iis/iisstandards/vaccines.asp?rpt=tradename>. **4.** Medicare NCCI 2020 Coding Policy Manual. Chapter XI Medicine evaluation and management services CPT codes 90000-99999 for national correct coding initiative policy manual for Medicare services. Last Revised January 1, 2020. Accessed October 30, 2020. https://www.supercoder.com/exclusives/cci-policymanual/Chapter11_CPTCodes90000-99999_Final_11_12_19.pdf.

VAXELIS is a trademark of MSP Vaccine Company. Brands mentioned are trademarks of their respective owners.

Copyright © 2021 MSP Vaccine Company. All Rights Reserved.
US-VAX-00055 11/20



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VAXELIS safely and effectively. See full prescribing information for VAXELIS.

VAXELIS™ (Diphtheria and Tetanus Toxoids and Acellular Pertussis, Inactivated Poliovirus, Haemophilus b Conjugate and Hepatitis B Vaccine)

Suspension for Intramuscular Injection

Initial U.S. Approval: 2018

-----RECENT MAJOR CHANGES-----

Dosage and Administration (2.2) 09/2020

-----INDICATIONS AND USAGE-----

VAXELIS is a vaccine indicated for active immunization to prevent diphtheria, tetanus, pertussis, poliomyelitis, hepatitis B, and invasive disease due to *Haemophilus influenzae* type b. VAXELIS is approved for use as a 3-dose series in children from 6 weeks through 4 years of age (prior to the 5th birthday). (1)

-----DOSAGE AND ADMINISTRATION-----

The 3-dose immunization series consists of a 0.5 mL intramuscular injection, administered at 2, 4, and 6 months of age. (2.1)

-----DOSAGE FORMS AND STRENGTHS-----

Suspension for injection (0.5 mL dose) available in single-dose vials and prefilled syringes. (3)

-----CONTRAINDICATIONS-----

- Severe allergic reaction (e.g., anaphylaxis) to a previous dose of VAXELIS, any ingredient of VAXELIS, or any other diphtheria toxoid, tetanus toxoid, pertussis-containing vaccine, inactivated poliovirus vaccine, hepatitis B vaccine, or *Haemophilus influenzae* type b vaccine. (4.1)
- Encephalopathy within 7 days of a previous pertussis-containing vaccine with no other identifiable cause. (4.2)
- Progressive neurologic disorder until a treatment regimen has been established and the condition has stabilized. (4.3)

-----WARNINGS AND PRECAUTIONS-----

- Carefully consider benefits and risks before administering VAXELIS to persons with a history of:
 - fever $\geq 40.5^{\circ}\text{C}$ ($\geq 105^{\circ}\text{F}$), hypotonic-hyporesponsive episode (HHE) or persistent, inconsolable crying lasting ≥ 3 hours within 48 hours after a previous pertussis-containing vaccine. (5.2)
 - seizures within 3 days after a previous pertussis-containing vaccine. (5.2)
- If Guillain-Barré syndrome occurred within 6 weeks of receipt of a prior vaccine containing tetanus toxoid, the risk for Guillain-Barré syndrome may be increased following VAXELIS. (5.3)
- Apnea following intramuscular vaccination has been observed in some infants born prematurely. The decision about when to administer an intramuscular vaccine, including VAXELIS, to an infant born prematurely should be based on consideration of the individual infant's medical status and the potential benefits and possible risks of vaccination. (5.5)
- Urine antigen detection may not have definitive diagnostic value in suspected *H. influenzae* type b disease following vaccination with VAXELIS. (5.7) (7.1)

-----ADVERSE REACTIONS-----

The solicited adverse reactions following any dose were irritability ($\geq 55\%$), crying ($\geq 45\%$), injection site pain ($\geq 44\%$), somnolence ($\geq 40\%$), injection site erythema ($\geq 25\%$), decreased appetite ($\geq 23\%$), fever $\geq 38.0^{\circ}\text{C}$ ($\geq 19\%$), injection site swelling ($\geq 18\%$), and vomiting ($\geq 9\%$). (6)

To report SUSPECTED ADVERSE REACTIONS, contact Sanofi Pasteur Inc., at 1-800-822-2463 (1-800-VACCINE) or VAERS at 1-800-822-7967 and <http://vaers.hhs.gov>.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 10/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Vaccination Schedule
- 2.2 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 4.1 Hypersensitivity
- 4.2 Encephalopathy
- 4.3 Progressive Neurologic Disorder

5 WARNINGS AND PRECAUTIONS

- 5.1 Management of Acute Allergic Reactions
- 5.2 Adverse Reactions Following Prior Pertussis Vaccination
- 5.3 Guillain-Barré Syndrome and Brachial Neuritis
- 5.4 Altered Immunocompetence
- 5.5 Apnea in Premature Infants
- 5.6 Limitations of Vaccine Effectiveness
- 5.7 Interference with Laboratory Tests

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Interference with Laboratory Tests

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Effectiveness of VAXELIS
- 14.2 Immunogenicity
- 14.3 Concomitantly Administered Vaccines

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

VAXELIS™ is a vaccine indicated for active immunization to prevent diphtheria, tetanus, pertussis, poliomyelitis, hepatitis B, and invasive disease due to *Haemophilus influenzae* (*H. influenzae*) type b. VAXELIS is approved for use as a 3-dose series in children 6 weeks through 4 years of age (prior to the 5th birthday).

2 DOSAGE AND ADMINISTRATION

For intramuscular use only.

2.1 Vaccination Schedule

VAXELIS is to be administered as a 3-dose series at 2, 4, and 6 months of age. The first dose may be given as early as 6 weeks of age. Three doses of VAXELIS constitute a primary immunization course against diphtheria, tetanus, *H. influenzae* type b invasive disease and poliomyelitis.

VAXELIS may be used to complete the hepatitis B immunization series.

A 3-dose series of VAXELIS does not constitute a primary immunization series against pertussis; an additional dose of pertussis-containing vaccine is needed to complete the primary series. [See *Pertussis Vaccination Following VAXELIS*.]

Pertussis Vaccination following VAXELIS

VAXELIS, Pentacel® [(Diphtheria and Tetanus Toxoids and Acellular Pertussis Adsorbed, Inactivated Poliovirus and Haemophilus b Conjugate (Tetanus Toxoid Conjugate) Vaccine): DTaP-IPV/Hib], Quadracel® [(Diphtheria and Tetanus Toxoids and Acellular Pertussis Adsorbed and Inactivated Poliovirus Vaccine): DTaP-IPV] and DAPTACEL® [(Diphtheria and Tetanus Toxoids and Acellular Pertussis Vaccine Adsorbed): DTaP] contain the same pertussis antigens manufactured by the same process. Children who have received a 3-dose series of VAXELIS should complete the primary and pertussis vaccination series with Pentacel, Quadracel or DAPTACEL according to the respective prescribing information in the approved package inserts. [See *ADVERSE REACTIONS (6.1) AND CLINICAL STUDIES (14)*.]

Administration of VAXELIS following previous doses of other DTaP-containing Vaccines

VAXELIS may be used to complete the first 3 doses of the 5-dose DTaP series in infants and children who have received 1 or 2 doses of Pentacel or DAPTACEL and are also scheduled to receive the other antigens in VAXELIS. Data are not available on the safety and immunogenicity of such mixed sequences.

Data are not available on the safety and effectiveness of using VAXELIS following 1 or 2 doses of a DTaP vaccine from a different manufacturer.

Administration of VAXELIS following previous doses of any Hepatitis B Vaccine

A 3-dose series of VAXELIS may be administered to infants born to HBsAg-negative mothers, and who have received a dose of any hepatitis B vaccine, prior to or at 1 month of age. [See *ADVERSE REACTIONS (6.1) AND CLINICAL STUDIES (14)*.]

VAXELIS may be used to complete the hepatitis B vaccination series following 1 or 2 doses of other hepatitis B vaccines, in infants and children born of HBsAg-negative mothers and who are also scheduled to receive the other antigens in VAXELIS. However, data are not available on the safety and effectiveness of VAXELIS in such infants and children.

Administration of VAXELIS following previous doses of Inactivated Polio Vaccine (IPV)

VAXELIS may be administered to infants and children who have received 1 or 2 doses of IPV and are also scheduled to receive the other antigens in VAXELIS. However, data are not available on the safety and effectiveness of VAXELIS in such infants and children.

Administration of VAXELIS following previous doses of Haemophilus b Conjugate Vaccines

VAXELIS may be administered to infants and children who have received 1 or 2 doses of *H. influenzae* type b Conjugate Vaccine and are also scheduled to receive the other antigens in VAXELIS. However, data are not available on the safety and effectiveness of VAXELIS in such infants and children.

2.2 Administration

Just before use, shake the vial or syringe until a uniform, white, cloudy suspension results.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. If either of these conditions exist, the product should not be administered.

Administer a single 0.5 mL dose of VAXELIS intramuscularly.

In infants younger than 1 year, the anterolateral aspect of the thigh is the preferred site of injection. The vaccine should not be injected into the gluteal area.

VAXELIS should not be combined through reconstitution or mixed with any other vaccine. Discard unused portion.

3 DOSAGE FORMS AND STRENGTHS

VAXELIS is a suspension for injection available in 0.5 mL single-dose vials and prefilled syringes. [See *HOW SUPPLIED/STORAGE AND HANDLING (16)*.]

4 CONTRAINDICATIONS

4.1 Hypersensitivity

Do not administer VAXELIS to anyone with a history of a severe allergic reaction (e.g., anaphylaxis) to a previous dose of VAXELIS, any ingredient of VAXELIS, or any other diphtheria toxoid, tetanus toxoid, pertussis-containing vaccine, inactivated poliovirus vaccine, hepatitis B vaccine, or *H. influenzae* type b vaccine [See *DESCRIPTION (11)*.]

4.2 Encephalopathy

Do not administer VAXELIS to anyone with a history of encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures) within 7 days of a previous dose of a pertussis-containing vaccine, that is not attributable to another identifiable cause.

4.3 Progressive Neurologic Disorder

Do not administer VAXELIS to anyone with a history of progressive neurologic disorder, including infantile spasms, uncontrolled epilepsy, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized.

5 WARNINGS AND PRECAUTIONS

5.1 Management of Acute Allergic Reactions

Epinephrine hydrochloride solution (1:1,000) and other appropriate agents and equipment must be available for immediate use in case an anaphylactic or acute hypersensitivity reaction occurs.

5.2 Adverse Reactions Following Prior Pertussis Vaccination

If any of the following events occur after administration of a pertussis vaccine, the decision to administer VAXELIS should be based on careful consideration of potential benefits and possible risks.

- Temperature of $\geq 40.5^{\circ}\text{C}$ ($\geq 105^{\circ}\text{F}$) within 48 hours, not attributable to another identifiable cause.
- Collapse or shock-like state (hypotonic-hyporesponsive episode [HHE]) within 48 hours.
- Persistent, inconsolable crying lasting ≥ 3 hours within 48 hours.
- Seizures with or without fever within 3 days.

5.3 Guillain-Barré Syndrome and Brachial Neuritis

A review by the Institute of Medicine (IOM) found evidence for a causal relation between tetanus toxoid and both brachial neuritis and Guillain-Barré syndrome. If Guillain-Barré syndrome occurred within 6 weeks of receipt of a prior vaccine containing tetanus toxoid, the risk for Guillain-Barré syndrome may be increased following VAXELIS. (1)

5.4 Altered Immunocompetence

If VAXELIS is administered to immunocompromised persons, including persons receiving immunosuppressive therapy, the expected immune response may not be obtained.

5.5 Apnea in Premature Infants

Apnea following intramuscular vaccination has been observed in some infants born prematurely. The decision about when to administer an intramuscular vaccine, including VAXELIS, to an infant born prematurely should be based on consideration of the infant's medical status and the potential benefits and possible risks of vaccination.

5.6 Limitations of Vaccine Effectiveness

Vaccination with VAXELIS may not protect all individuals.

5.7 Interference with Laboratory Tests

Urine antigen detection may not have definitive diagnostic value in suspected *H. influenzae* type b disease following vaccination with VAXELIS. [See *DRUG INTERACTIONS (7.1)*.]

6 ADVERSE REACTIONS

Rates of adverse reactions varied by number of doses of VAXELIS received. The solicited adverse reactions 0-5 days following any dose were irritability ($\geq 55\%$), crying ($\geq 45\%$), injection site pain ($\geq 44\%$), somnolence ($\geq 40\%$), injection site erythema ($\geq 25\%$), decreased appetite ($\geq 23\%$), fever $\geq 38.0^{\circ}\text{C}$ ($\geq 19\%$), injection site swelling ($\geq 18\%$), and vomiting ($\geq 9\%$).

6.1 Clinical Trials Experience

Because clinical trials are conducted under varying conditions, adverse reaction rates observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trials of another vaccine and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to vaccine use and for approximating rates of those events.

The safety of VAXELIS was evaluated in 6 clinical studies, in which a total of 5,251 infants 43 to 99 days of age at enrollment received at least 1 dose of VAXELIS. Two of these (study 005 and 006) were controlled clinical studies conducted in the US, in which a total of 3,380 infants 46 to 89 days of age at enrollment received at least 1 dose of VAXELIS. The vaccination schedules of VAXELIS, Control vaccines, and concomitantly administered vaccines used in these studies are provided in Table 1. At 15 months of age, participants in Study 005 received a dose of DAPTACEL and a *H. influenzae* type b conjugate vaccine, whereas participants in Study 006 received a dose of Pentacel. In a non-US study, 294 children received a dose of VAXELIS at 15 months of age.

Across the 2 studies conducted in the US, among all randomized participants (3,392 in the VAXELIS group and 889 in the Control group), 52.6% were male and 47.4% were female. The race distribution was as follows: 71.7% were White, 11.0% were Black, 4.5% were American Indian or Alaska Native, 3.5% were Asian, and 9.3% were of other racial groups. Most participants (81.8%) were of non-Hispanic or Latino ethnicity. The racial/ethnic distribution of participants who received VAXELIS and Control vaccines was similar.

Table 1: Clinical Safety Studies with VAXELIS in the US: Vaccination Schedules

Study	Vaccine	Concomitantly Administered Vaccines
005*	VAXELIS at 2, 4, 6 months and DAPTACEL + PedvaxHIB® at 15 months	RotaTeq® at 2, 4, and 6 months Pevnar 13® at 2, 4, 6, and 15 months
	Control group vaccines: Pentacel at 2, 4, 6 months and RECOMBIVAX HB® at 2 and 6 months DAPTACEL+ ActHIB® at 15 months	RotaTeq at 2, 4, and 6 months Pevnar 13 at 2, 4, 6, and 15 months
006*	VAXELIS at 2, 4, 6 months and Pentacel at 15 months	RotaTeq at 2, 4, and 6 months Pevnar 13 at 2, 4, 6, and 15 months
	Control group vaccines: Pentacel at 2, 4, 6, and 15 months RECOMBIVAX HB at 2 and 6 months	RotaTeq at 2, 4, and 6 months Pevnar 13 at 2, 4, 6, and 15 months

Pevnar 13 (Pneumococcal 13-valent Conjugate Vaccine [Diphtheria CRM197 Protein])

RotaTeq (Rotavirus Vaccine, Live, Oral, Pentavalent)

PedvaxHIB [Haemophilus b Conjugate Vaccine (Meningococcal Protein Conjugate)]

RECOMBIVAX HB (Hepatitis B Vaccine [Recombinant])

* The first dose of Hepatitis B vaccine was administered prior to study initiation (prior to or at 1 month of age).

Solicited Adverse Reactions

Information on solicited adverse events was recorded daily by parents or guardians on vaccination report cards. The incidence and severity of solicited injection site and systemic adverse reactions (i.e., vaccine-related adverse events) that occurred within 5 days following each dose of VAXELIS or Control vaccines at 2, 4, and 6 months of age in studies 005 and 006 are shown in Table 2.

Table 2: Percentage of Infants with Solicited Adverse Reactions Occurring within 5 days Following VAXELIS or Control Vaccines Administered Concomitantly at Separate Sites with Prevnar 13 and RotaTeq in Studies 005 and 006

		VAXELIS + Prevnar 13 + RotaTeq			Pentacel + RECOMBIVAX HB + Prevnar 13 + RotaTeq		
		Dose 1 (N=3,370) (%)	Dose 2 (N=3,221) (%)	Dose 3 (N=3,134) (%)	Dose 1 (N=880) (%)	Dose 2 (N=849) (%)	Dose 3 (N=825) (%)
Injection Site Adverse Reactions		VAXELIS site			Pentacel or RECOMBIVAX HB site		
Injection site erythema	Any	25.8	31.8	31.8	25.0	25.8	30.9
	≥2.5 cm	0.9	1.0	1.3	1.1	1.1	1.2
	>5.0 cm	0.0	0.1	0.2	0.3	0.2	0.1
Injection site pain*	Any	53.3	49.0	44.9	55.8	43.7	44.4
	Moderate or severe	16.3	14.1	12.5	19.1	11.3	10.8
	Severe	2.8	2.5	2.0	3.2	1.9	1.3
Injection site swelling	Any	18.9	22.8	23.4	20.8	20.4	22.9
	≥2.5 cm	2.5	1.6	1.7	2.7	1.3	0.8
	>5.0 cm	0.2	0.2	0.2	0.3	0.1	0.0
Systemic Adverse Reactions							
Fever	≥38°C	19.2	29.0	29.3	14.6	18.0	17.8
	≥38.5°C	5.3	11.5	13.2	3.4	6.5	8.1
	≥39.5°C	0.2	0.7	1.5	0.1	0.2	0.9
Crying	Any	52.0	49.5	45.1	50.6	47.0	40.6
	>1 hour	18.6	19.8	16.7	20.6	16.8	14.1
	>3 hours	3.6	3.8	3.4	4.4	4.0	2.9
Decreased Appetite[†]	Any	28.9	24.2	23.2	25.8	20.5	20.1
	Moderate or severe	7.0	5.5	4.8	6.8	3.9	5.0
	Severe	0.5	0.5	0.5	0.6	0.2	0.0
Irritability[‡]	Any	61.8	58.9	55.2	61.7	56.3	51.6
	Moderate or severe	24.6	23.4	20.1	25.7	19.2	16.8
	Severe	2.5	3.8	2.9	2.2	2.7	2.2

		VAXELIS + Prevnar 13 + RotaTeq			Pentacel + RECOMBIVAX HB + Prevnar 13 + RotaTeq		
		Dose 1 (N=3,370) (%)	Dose 2 (N=3,221) (%)	Dose 3 (N=3,134) (%)	Dose 1 (N=880) (%)	Dose 2 (N=849) (%)	Dose 3 (N=825) (%)
Somnolence[§]	Any	56.3	47.8	40.8	55.2	44.1	38.8
	Moderate or severe	15.0	11.5	8.5	14.5	9.4	8.2
	Severe	1.5	1.1	1.0	1.7	0.6	1.1
Vomiting[¶]	Any	13.1	11.5	9.5	11.3	9.7	6.9
	Moderate or severe	3.5	2.6	2.1	2.8	3.1	1.0
	Severe	0.4	0.2	0.1	0.5	0.6	0.1

N = Number of vaccinated participants with safety follow-up.

* Moderate: cries and protests when injection site is touched; Severe: cries when injected limb is moved or the movement of the injected limb is reduced.

† Moderate: missed 1 or 2 feeds/meals completely; Severe: refuses ≥ 3 feeds or refuses most feeds.

‡ Moderate: requiring increased attention; Severe: inconsolable.

§ Moderate: not interested in surroundings or did not wake up for a meal; Severe: Sleeping most of the time or difficult to wake up.

¶ Moderate: 2-5 episodes per 24 hours; Severe: ≥ 6 episodes per 24 hours or requiring parenteral hydration.

A subject with the same adverse reactions at both the Pentacel and RECOMBIVAX HB injection site, was counted once and was classified according to the highest intensity grading.

Fever is based upon actual temperatures recorded with no adjustments due to the measurement route.

Following Doses 1-3 combined, the proportion of temperature measurements that were taken by rectal, axillary, or other routes were 91.7%, 8.1%, and 0% respectively, for VAXELIS group, and 90.3%, 9.7%, and 0%, respectively, for Pentacel + RECOMBIVAX HB vaccines group.

Non-fatal Serious Adverse Events

Across Studies 005 and 006, within 30 days following any infant dose vaccination, 68 participants (2.0%) who received VAXELIS and concomitant vaccines versus 19 participants (2.2%) who received Control and concomitant vaccines experienced a serious adverse event. Of these, a vaccine-related SAE was reported for no participants in the Control vaccines group and for 4 participants (0.1%) in the VAXELIS group:

- 3 of these 4 experienced pyrexia 1 to 2 days following the first study vaccinations; and
- 1 of these 4 experienced an apparent life-threatening event (vomiting followed by pallor and lethargy) on the day of the first study vaccinations, and again 2 days later.

Deaths

In the 2 US studies, death was reported in 6 participants (0.2%) who received VAXELIS and in 1 participant (0.1%) who received Pentacel + RECOMBIVAX HB vaccines; none were assessed as vaccine related. Causes of death among infants who received VAXELIS were asphyxia, hydrocephalus, unknown cause, sepsis and 2 cases of Sudden Infant Death Syndrome (occurring 1, 2, 10, 42, 44 and 49 days post-vaccination, respectively). Across all 6 clinical studies, there were no deaths assessed as related to VAXELIS.

6.2 Postmarketing Experience

The following adverse events have been reported during post-marketing use of VAXELIS or other vaccines containing the antigens of VAXELIS. These adverse events are included based on a suspected causal connection to VAXELIS or the components of DAPTACEL[®] (Diphtheria and Tetanus Toxoids and Acellular Pertussis Vaccine Adsorbed), IPOL[®] (Poliovirus Vaccine Inactivated), COMVAX[®] [Haemophilus b Conjugate (Meningococcal Protein Conjugate) and Hepatitis B (Recombinant) Vaccine].

Because these events are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to vaccination.

- ***Immune System Disorders***
Hypersensitivity (such as rash, urticaria, dyspnea, erythema multiforme), anaphylactic reaction (such as urticaria, angioedema, edema, face edema, shock).
- ***General Disorders and Administration Site Conditions***
Extensive swelling of injected limb (including swelling that involves adjacent joints).
- ***Nervous System***
Seizure, febrile seizure, hypotonic-hyporesponsive episode (HHE).

7 DRUG INTERACTIONS

7.1 Interference with Laboratory Tests

Sensitive tests (e.g., Latex Agglutination kits) have detected vaccine-derived polyribosylribitol phosphate (PRP) in the urine of vaccinees for at least 30 days following vaccination with PedvaxHIB [Haemophilus b Conjugate Vaccine (Meningococcal Protein Conjugate)]. (2) Therefore, urine antigen detection may not have definite diagnostic value in suspected *H. influenzae* type b disease following vaccination with VAXELIS. [See **WARNINGS AND PRECAUTIONS** (5.7).]

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

VAXELIS is not approved for use in individuals 5 years of age and older. No human or animal data are available to assess vaccine-associated risks in pregnancy.

8.2 Lactation

VAXELIS is not approved for use in individuals 5 years of age and older. No human or animal data are available to assess the impact of VAXELIS on milk production, its presence in breast milk, or its effects on the breastfed infant.

8.4 Pediatric Use

The safety of VAXELIS has been established in the age group 6 weeks through 15 months, and the effectiveness of VAXELIS was established in the age group 6 weeks through 6 months on the basis of clinical studies. [See *ADVERSE REACTIONS (6.1) AND CLINICAL STUDIES (14).*]

The safety and effectiveness of VAXELIS in older children through 4 years of age are supported by evidence in younger children. The safety and effectiveness of VAXELIS in infants less than 6 weeks of age and in children and adolescents 5 through 17 years of age have not been established.

11 DESCRIPTION

VAXELIS (Diphtheria and Tetanus Toxoids and Acellular Pertussis, Inactivated Poliovirus, Haemophilus b Conjugate and Hepatitis B Vaccine) is a sterile suspension for intramuscular injection.

Each 0.5 mL dose is formulated to contain 15 Lf diphtheria toxoid, 5 Lf tetanus toxoid, acellular pertussis antigens [20 mcg detoxified pertussis toxin (PT), 20 mcg filamentous hemagglutinin (FHA), 3 mcg pertactin (PRN), 5 mcg fimbriae types 2 and 3 (FIM)], inactivated polioviruses [29 D-antigen units (DU) Type 1 (Mahoney), 7 DU Type 2 (MEF-1), 26 DU Type 3 (Saukett)], 3 mcg polyribosylribitol phosphate (PRP) of *H. influenzae* type b covalently bound to 50 mcg of the outer membrane protein complex (OMPC) of *Neisseria meningitidis* serogroup B, and 10 mcg hepatitis B surface antigen (HBsAg). Each 0.5 mL dose contains 319 mcg aluminum from aluminum salts used as adjuvants.

Other ingredients per 0.5 mL dose include <0.0056% polysorbate 80 and the following residuals from the manufacturing process: ≤14 mcg formaldehyde, ≤50 ng glutaraldehyde, ≤50 ng bovine serum albumin, <5 ng of neomycin, <200 ng streptomycin sulfate, <25 ng polymyxin B sulfate, ≤0.125 µg ammonium thiocyanate and ≤0.1 mcg yeast protein (maximum 1% relative to HBsAg protein).

Corynebacterium diphtheriae is grown in modified Mueller's growth medium. (3) After purification by ammonium sulfate fractionation, the diphtheria toxin is detoxified with formaldehyde and diafiltered.

Clostridium tetani is grown in modified Mueller-Miller casamino acid medium without beef heart infusion. (4) Tetanus toxin is detoxified with formaldehyde and purified by ammonium sulfate fractionation and diafiltration. Diphtheria and tetanus toxoids are individually adsorbed onto aluminum phosphate.

The acellular pertussis vaccine antigens are produced from *Bordetella pertussis* cultures grown in Stainer-Scholte medium (5) modified by the addition of casamino acids and dimethyl-beta-cyclodextrin. PT, FHA and PRN are isolated separately from the supernatant culture medium. FIM are extracted and copurified from the bacterial cells. The pertussis antigens are purified by sequential filtration, salt-precipitation, ultrafiltration and chromatography. PT is detoxified with glutaraldehyde. FHA is treated with formaldehyde and the residual aldehydes are removed by ultrafiltration. The individual antigens are adsorbed separately onto aluminum phosphate.

The Type 1, Type 2, and Type 3 polioviruses are individually grown in Vero cells. The viral harvests are concentrated and purified, then inactivated with formaldehyde to produce

monovalent suspensions of each serotype. Specified quantities of monovalent suspensions of each serotype are mixed to produce the trivalent poliovirus concentrate.

The HBsAg antigen is harvested and purified from fermentation cultures of a recombinant strain of the yeast *Saccharomyces cerevisiae* containing the gene for the *adw* subtype of HBsAg. The recombinant *Saccharomyces cerevisiae* is grown in a fermentation medium which consists of an extract of yeast, soy peptone, dextrose, amino acids, and mineral salts. The HBsAg protein is released from the yeast cells by cell disruption and purified by a series of physical and chemical methods which includes ion and hydrophobic chromatography, and diafiltration. The purified protein is treated in phosphate buffer with formaldehyde and then co-precipitated with alum (potassium aluminum sulfate) to form bulk vaccine adjuvanted with amorphous aluminum hydroxyphosphate sulfate.

The purified PRP of *H. influenzae* type b (Haemophilus b, Ross strain) is conjugated to an OMPC of the B11 strain of *N. meningitidis* serogroup B. *H. influenzae* type b is grown in a fermentation medium which includes an extract of yeast, nicotinamide adenine dinucleotide, hemin chloride, soy peptone, dextrose, and mineral salts. The PRP is purified from the culture broth by purification procedures which include ethanol fractionation, enzyme digestion, phenol extraction and diafiltration. *N. meningitidis* serogroup B is grown in a fermentation medium which includes an extract of yeast, amino acids and mineral salts. The OMPC is purified by detergent extraction, ultracentrifugation, diafiltration and sterile filtration. PRP is conjugated to OMPC by chemical coupling and the PRP-OMPC is then adsorbed onto an amorphous aluminum hydroxyphosphate sulfate adjuvant.

The adsorbed diphtheria, tetanus, and acellular pertussis antigens are combined with aluminum phosphate (as adjuvant) and water for injection into an intermediate concentrate. The individual HBsAg and PRP-OMPC adjuvanted bulks are added followed by the trivalent poliovirus concentrate, to produce VAXELIS.

Both diphtheria and tetanus toxoids induce at least 2 neutralizing units per mL of serum in the guinea pig potency test. The potency of the acellular pertussis antigens is evaluated by the antibody response of immunized mice to detoxified PT, FHA, PRN and FIM as measured by enzyme-linked immunosorbent assay (ELISA). The immunogenicity of the inactivated polioviruses is evaluated by the antibody response in rats measured by virus neutralization. The potency of the HBsAg component is measured relative to a standard by an *in vitro* immunoassay. The potency of the PRP-OMPC component is measured by quantitating the polysaccharide concentration using an HPLC method.

VAXELIS does not contain a preservative. The vial stopper, syringe plunger stopper, and syringe tip cap are not made with natural rubber latex.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Diphtheria

Diphtheria is an acute toxin-mediated disease caused by toxigenic strains of *C. diphtheriae*. Protection against disease is due to the development of neutralizing antibodies to diphtheria toxin. A serum diphtheria antitoxin level of 0.01 IU/mL is the lowest level giving some degree of

protection. Antitoxin levels of ≥ 0.1 IU/mL are generally regarded as protective. (6) Levels of 1.0 IU/mL have been associated with long-term protection. (7)

Tetanus

Tetanus is an acute disease caused by an extremely potent neurotoxin produced by *C. tetani*. Protection against disease is due to the development of neutralizing antibodies to tetanus toxin. A serum tetanus antitoxin level of ≥ 0.01 IU/mL, measured by neutralization assay is considered the minimum protective level. (6) (8) A tetanus antitoxoid level ≥ 0.1 IU/mL as measured by the ELISA used in clinical studies of VAXELIS is considered protective.

Pertussis

Pertussis (whooping cough) is a respiratory disease caused by *B. pertussis*. This Gram-negative coccobacillus produces a variety of biologically active components, though their role in either the pathogenesis of, or immunity to, pertussis has not been clearly defined.

Poliomyelitis

Polioviruses, of which there are 3 serotypes (Types 1, 2, and 3), are enteroviruses. The presence of poliovirus type-specific neutralizing antibodies has been correlated with protection against poliomyelitis. (9)

Hepatitis B

Hepatitis B virus is one of several hepatitis viruses that cause systemic infection, with major pathology in the liver. Antibody concentrations of ≥ 10 mIU/mL against HBsAg correlate with protection against hepatitis B virus infection.

Haemophilus influenzae type b Invasive Disease

H. influenzae type b can cause invasive disease such as meningitis and sepsis. Anti-PRP antibody has been shown to correlate with protection against invasive disease due to *H. influenzae* type b.

Based on data from passive antibody studies (10) and an efficacy study with *H. influenzae* type b polysaccharide vaccine in Finland, (11) a post-vaccination anti-PRP level of ≥ 0.15 mcg/mL is considered a minimal protective level. Data from an efficacy study with *H. influenzae* type b polysaccharide vaccine in Finland indicate that an anti-PRP level of ≥ 1.0 mcg/mL 3 weeks after vaccination predicts protection through a subsequent 1-year period. (11) (12) These levels have been used to evaluate the effectiveness of *H. influenzae* type b conjugate vaccines, including the PRP-OMPC component of VAXELIS.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

VAXELIS has not been evaluated for carcinogenic or mutagenic potential or impairment of fertility.

14 CLINICAL STUDIES

14.1 Effectiveness of VAXELIS

The effectiveness of VAXELIS is based on the immunogenicity of the individual antigens compared to US licensed vaccines. Serological correlates of protection exist for diphtheria,

tetanus, hepatitis B, poliomyelitis, and invasive disease due to *H. influenzae* type b. The effectiveness against pertussis is based upon the pertussis immune responses following 3 doses of VAXELIS compared to 3 doses of Pentacel, as well as the pertussis immune responses following a subsequent dose of DAPTACEL in the same 2 groups of children. VAXELIS, Pentacel and DAPTACEL contain the same pertussis antigens, manufactured by the same processes.

14.2 Immunogenicity

In the US Study 005 (Table 1), infants were randomized to receive 3 doses of VAXELIS at 2, 4, and 6 months of age and DAPTACEL and PedvaxHIB at 15 months of age, or Control group vaccines (3 doses of Pentacel vaccine at 2, 4, and 6 months of age + RECOMBIVAX HB at 2 and 6 months of age and DAPTACEL and ActHIB at 15 months of age). All subjects received concomitant vaccines: RotaTeq at 2, 4 and 6 months and Prevnar 13 at 2, 4, 6, and 15 months of age. [See *ADVERSE REACTIONS (6.1)*.] All infants had received a dose of hepatitis B vaccine prior to study initiation, prior to or at one month of age. Among all randomized participants, 53.0% were male and 47.0% were female. Most (79.2%) participants were White, 14.1% were Black and 5.2% were multi-racial. Most (91.4%) participants were of non-Hispanic or non-Latin ethnicity.

Antibody responses to diphtheria, tetanus, pertussis (PT, FHA, PRN and FIM), poliovirus types 1, 2 and 3, hepatitis B and *H. influenzae* type b antigens were measured in sera obtained one month following the third dose of VAXELIS or Pentacel + RECOMBIVAX HB vaccines. VAXELIS was non-inferior to Pentacel + RECOMBIVAX HB administered concomitantly at separate sites, as demonstrated by the proportions of participants achieving seroprotective levels of antibodies to diphtheria, tetanus, poliovirus, hepatitis B and PRP antigens, and pertussis vaccine response rates and GMCs (except FHA), following 3 doses of the vaccine. See Table 3.

To complete the 4-dose pertussis primary vaccination series, participants in both groups received DAPTACEL at 15 months of age and were evaluated for immune responses to pertussis antigens one month later. The non-inferiority criteria for vaccine response rates and GMCs for all pertussis antigens were met following the fourth dose.

Table 3: Antibody Responses One Month Following Dose 3 of VAXELIS or Control Vaccines Administered Concomitantly with Prevnar 13 and RotaTeq in Study 005

	VAXELIS + Prevnar 13 + RotaTeq (N=688 - 810)	Pentacel + RECOMBIVAX HB + Prevnar 13 + RotaTeq (N=353 - 400)
Anti-Diphtheria Toxoid % ≥ 0.1 IU/mL	82.4*	86.3
Anti-Tetanus Toxoid % ≥ 0.1 IU/mL	99.9 [†]	99.5
Anti-PT % vaccine response [‡] GMC	98.1* 109.6 [§]	98.5 85.4
Anti-FHA % vaccine response [‡] GMC	87.3* 46.6 [¶]	92.0 72.3
Anti-PRN % vaccine response [‡] GMC	79.3* 55.8 [§]	82.0 66.8
Anti-FIM % vaccine response [‡] GMC	90.2* 235.9 [§]	86.2 184.4
Anti-Poliovirus Type 1 % $\geq 1:8$ dilution	100.0 [†]	98.2
Anti-Poliovirus Type 2 % $\geq 1:8$ dilution	100.0 [†]	99.7
Anti-Poliovirus Type 3 % $\geq 1:8$ dilution	100.0 [†]	99.8
Anti-PRP % ≥ 0.15 $\mu\text{g/mL}$ % ≥ 1.0 $\mu\text{g/mL}$	97.3 [†] 85.0*	92.4 75.3
Anti-HBsAg % ≥ 10 mIU/mL	99.4*	98.6

N = The number of participants with available data.

* Non-inferiority criterion met (lower bound of 2-sided 95% CI for the difference [VAXELIS group minus Control vaccines group] was $>-10\%$).

[†] Non-inferiority criterion met (lower bound of 2-sided 95% CI for the difference [VAXELIS group minus Control vaccines group] was $>-5\%$).

- ‡ Vaccine response = if pre-vaccination antibody concentration was $<4 \times$ lower limit of quantitation [LLOQ], then the post-vaccination antibody concentration was $\geq 4 \times$ LLOQ; if pre-vaccination antibody concentration was $\geq 4 \times$ LLOQ, then the post-vaccination antibody concentration was \geq pre-vaccination levels (pre-Dose 1).
- § Non-inferiority criterion met (lower bound of 2-sided 95% CI for the GMC ratio [VAXELIS group/Control vaccines group] was >0.67).
- ¶ Non-inferiority criterion not met for anti-FHA GMC (lower bound of 2-sided 95% CI for the GMC ratio [VAXELIS group/Control vaccines group] was 0.59 which is below the non-inferiority criterion >0.67).

Study 006 (Table 1) was a lot consistency study conducted in the US, where infants were randomized to receive 3 doses of VAXELIS at 2, 4, and 6 months of age and Pentacel at 15 months of age (N=2,406), or control group vaccines (4 doses of Pentacel at 2, 4, 6, and 15 months of age + RECOMBIVAX HB at 2 and 6 months of age; N=402). All subjects received concomitant vaccines: RotaTeq at 2, 4 and 6 months and Prevnar 13 at 2, 4, 6, and 15 months of age. All infants had received a dose of hepatitis B vaccine prior to study initiation, from birth up to one month of age.

Antibody responses to diphtheria, tetanus, pertussis (PT, FHA, PRN and FIM), poliovirus types 1, 2 and 3, hepatitis B and *H. influenzae* type b antigens were measured in sera obtained one month following the third dose of VAXELIS or Pentacel + RECOMBIVAX HB. VAXELIS was non-inferior to Pentacel + RECOMBIVAX HB administered concomitantly at separate sites, as demonstrated by the proportions of participants achieving seroprotective levels of antibodies to diphtheria, tetanus, poliovirus, hepatitis B and PRP antigens, and pertussis vaccine response rates and GMCs, except for GMCs for FHA (lower bound of 2-sided 95% CI for GMC ratio [VAXELIS group/Control group vaccines] was 0.62, which was below the non-inferiority criterion >0.67).

To complete the 4-dose pertussis primary vaccination series, participants in both groups received Pentacel at 15 months of age and were evaluated for immune responses to pertussis antigens one month later. The non-inferiority criteria for antibody vaccine response rates and GMCs for all pertussis antigens were met following the fourth dose except for GMCs for PRN (lower bound of 2-sided 95% CI for GMC ratio [VAXELIS group/Control group vaccines] was 0.66, which was below the non-inferiority criterion >0.67).

14.3 Concomitantly Administered Vaccines

In Study 006 conducted in the US (Table 1), the immune responses to Prevnar 13 were measured one month after the third dose. Non-inferiority criteria were met for GMCs to 12 of the 13 serotype antigens in Prevnar 13 for participants who received VAXELIS relative to Control vaccines. For serotype 6B, the non-inferiority criterion was not met (lower bound of 2-sided 95% CI for GMC ratio [VAXELIS group/Control vaccines group] is 0.64, which is below the non-inferiority criterion >0.67).

15 REFERENCES

- 1 Stratton K, Ford A, Rusch E, Clayton EW, eds. Institute of Medicine (IOM). Adverse Effects of Vaccines: Evidence and Causality. Washington, DC: The National Academies Press. 2011.
- 2 Goepp JG, Hohenboken M, Almeida-Hill J, Santosham M. Persistent urinary antigen excretion in infants vaccinated with *Haemophilus influenzae* type b capsular polysaccharide conjugated with outer membrane protein from *Neisseria meningitidis*. *Pediatr Infect Dis J* 1992;11(1):2-5.
- 3 Stainer DW. Production of diphtheria toxin. In: Manclark CR, editor. Proceedings of an informal consultation on the World Health Organization requirements for diphtheria, tetanus, pertussis and combined vaccines. United States Public Health Service, Bethesda, MD. DHHS 91-1174; 1991. p. 7-11.
- 4 Mueller JH, Miller PA. Variable factors influencing the production of tetanus toxin. *J Bacteriol* 1954;67(3):271-7.
- 5 Stainer DW, Scholte MJ. A simple chemically defined medium for the production of phase I Bordetella pertussis. *J Gen Microbiol* 1971;63:211-20.
- 6 Department of Health and Human Services, Food and Drug Administration. Biological products; bacterial vaccines and toxoids; implementation of efficacy review; proposed rule. *Federal Register* 1985;50(240):51002-117.
- 7 Tiwari TSP, Wharton M. Diphtheria toxoid. In: Plotkin SA, Orenstein WA, and Offit PA, editors. *Vaccines*. 6th ed. Philadelphia, PA: WB Saunders; 2013:153-66.
- 8 Roper M, Wassilak SGF, Tiwari TSP, Orenstein WA. Tetanus toxoid. In: Plotkin SA, Orenstein WA, Offit PA, editors. *Vaccines*. 6th ed. Philadelphia, PA: WB Saunders; 2013. p. 746-72.
- 9 Sutter RW, et al. Defining surrogate serologic tests with respect to predicting protective vaccine efficacy: Poliovirus vaccination. In: Williams JC, et al. eds. *Combined vaccines and simultaneous administration. Current issues and perspectives*. New York, NY: The New York Academy of Sciences. 1995:289-99.
- 10 Robbins, J. B., et al: Quantitative measurement of 'natural' and immunization-induced *Haemophilus influenzae* type b capsular polysaccharide antibodies. *Pediatr Res* 1973;7(3):103-10.
- 11 Kayhty H, et al. The protective level of serum antibodies to the capsular polysaccharide of *Haemophilus influenzae* type b. *J Infect Dis* 1983;147:1100.
- 12 Anderson P. The protective level of serum antibodies to the capsular polysaccharide of *Haemophilus influenzae* type b. *J Infect Dis* 1984;149:1034.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Single-dose vial (NDC 63361-243-58) in packages of 10 vials (NDC 63361-243-10).

Single-dose, prefilled syringe with Luer lock connection and a tip cap, without needle, 0.5 mL (NDC 63361-243-88). Supplied as package of 10 (NDC 63361-243-15).

The vial stopper, syringe plunger stopper, and syringe tip cap are not made with natural rubber latex.

16.2 Storage and Handling

VAXELIS should be stored at 2°C to 8°C (36°F to 46°F). **Do not freeze.** Product which has been exposed to freezing should not be used. Protect from light. Do not use after expiration date shown on the label. Discard unused portion.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Inform the parent or guardian of the following:

- The potential benefits and risks of immunization with VAXELIS.
- The common adverse reactions that have occurred following administration of VAXELIS or other vaccines containing similar ingredients.
- Other adverse reactions can occur. Call healthcare provider with any adverse reactions of concern.

Provide the Vaccine Information Statements (VIS), which are required by the National Childhood Vaccine Injury Act of 1986.

Manufactured by:

Sanofi Pasteur Limited
Toronto Ontario Canada

for:

MSP Vaccine Company
Swiftwater PA 18370 USA

Distributed by:

Merck Sharp & Dohme Corp.
A subsidiary of **Merck & Co., Inc.**
Whitehouse Station NJ 08889 USA
and **Sanofi Pasteur Inc.**
Swiftwater PA 18370 USA

VAXELIS is a trademark of MSP Vaccine Company. The trademarks depicted herein are owned by their respective companies.

R6-1020 USA

Patient Information

VAXELIS™ (pronounced "vak-sel-lis")

(Diphtheria and Tetanus Toxoids and Acellular Pertussis, Inactivated Poliovirus, Haemophilus b Conjugate and Hepatitis B Vaccine)

Before your child gets VAXELIS, read this document and be sure you understand all of the information. Keep this document, you may need to read it again. If you have questions or side effects, ask your child's healthcare provider. This information does not take the place of talking about VAXELIS with your child's healthcare provider.

What is VAXELIS?

VAXELIS is a vaccine given to protect your child from getting diphtheria, tetanus (lockjaw), pertussis (whooping cough), polio, Hib (*Haemophilus influenzae* type b), and hepatitis B.

Your child cannot get any of these diseases from VAXELIS.

VAXELIS may not completely protect your child from these diseases.

Who should not get VAXELIS?

Your child should not get VAXELIS, if your child:

- is allergic to any of the ingredients.
- had an allergic reaction to any prior shot for diphtheria, tetanus, pertussis, polio, Hib, or hepatitis B.
- had a serious reaction affecting their brain and nervous system after a whooping cough shot.
- has a brain and nerve illness that is getting worse.

Before your child gets VAXELIS, tell your healthcare provider if your child:

- had problems with any shots for these diseases.
- is taking steroids, getting treatment for cancer, or has another problem that weakens the immune system.

How is VAXELIS given?

- VAXELIS is given to children from 6 weeks through 4 years of age (up to 5th birthday).
- Your child will need 3 shots:
 - one shot at 2 months old, **and**
 - one shot at 4 months old, **and**
 - one shot at 6 months old
- If your child misses a shot of VAXELIS, your healthcare provider may suggest a catch-up schedule.
- Your child may get VAXELIS at the same time they get other vaccines.

What are the most common side effects of VAXELIS?

- pain, redness, or swelling where the shot was given
- fever (100.4°F or higher)
- crying more than usual

- eating less than usual
- fussy more than usual
- sleepy more than usual
- throwing up

There may be other side effects that are not listed. If your child has any side effects that worry you or seem to get worse, tell your child's healthcare provider right away.

You may report any side effects directly to the Vaccine Adverse Event Reporting System (VAERS) at 1-800-822-7967 or <http://vaers.hhs.gov>, or contact Sanofi Pasteur Inc., at 1-800-822-2463 (1-800-VACCINE).

To learn more about VAXELIS, ask your healthcare provider. You can also find the Full Prescribing Information written for doctors at www.fda.gov/media/119465/download.

What is in VAXELIS?

- Active ingredients: inactivated bacteria of diphtheria, tetanus, pertussis, Hib, and inactivated hepatitis B and polio viruses. The bacteria and viruses in VAXELIS are not alive and do not cause disease.
- Other ingredients: aluminum salts, polysorbate 80, glutaraldehyde, formaldehyde, bovine serum albumin, neomycin, streptomycin, polymyxin B, ammonium thiocyanate, yeast protein, and water.
- VAXELIS does not have any preservatives in it.
- VAXELIS vial stopper, syringe plunger stopper, and syringe tip cap do not contain natural rubber latex.

Manufactured by: **Sanofi Pasteur Limited** Toronto Ontario Canada for: MSP Vaccine Company
Swiftwater PA 18370 USA.

Distributed by: **Merck Sharp & Dohme Corp.**, a subsidiary of **Merck & Co., Inc.** Whitehouse Station NJ
08889 USA, and by **Sanofi Pasteur Inc.** Swiftwater PA 18370 USA.

VAXELIS is a trademark of MSP Vaccine Company. The trademarks depicted herein are owned by their respective companies.

Initial Approval: 23 October 2020

R0-1020